

**Nikhil Verma, MD**  
**Hailey Merk, MSM, MMS, PA-C**  
**Lisa Raaf, MS, PA-C**  
**1611 W. Harrison, Suite #300**  
**Chicago, IL 60612**  
**Vermapa@rushortho.com**  
**Fax: 708-409-5179**  
[www.sportssurgerychicago.com](http://www.sportssurgerychicago.com)

MIDWEST  
**ORTHOPAEDICS**  
 at RUSH



## LATERJET OPEN ANTERIOR SHOULDER STABILIZATION

### Phase I MAXIMAL PROTECTION (0-4 Weeks Post-Op)

- Immobilization** for 4 weeks using sling.
- Elbow A/AAROM: flexion and extension.
- Protect anterior and posterior capsule from stretch, but begin passive ROM**
- Limit FE (supine forward elevation in the scapular plane) to 90 degrees
- Limit ER (external rotation) to neutral 30 degrees
- Do Not** perform Pendulums.
- Modalities (i.e. CryoCuff) PRN(as needed).
- Wrist and gripping excercises.
- Begin Deltoid/Cuff **isometrics**
- Removal of sling for showering: **maintain arm in sling position.**

### Phase II MODERATE PROTECTION (4-6 Weeks Post-Op)

- A/AAROM Limit FE (forward elevation in the scapular plane) to 140 degrees
- A/AAROM Limit ER (external rotation) to 45 degrees
- Progress from AAROM to AROM:
  - 1) Quality movement only-**avoid forcing** active motion with substitution patterns.
  - 2) Remember the effects of gravity on the limb, do gravity eliminated motions first ie. Supine elevation in the scapular plane.
- Deltoid isometrics.
- Elbow AROM
- Continue with wrist excercises
- Modalities PRN.
- Discontinue sling at 4-6 weeks.

### Phase III MINIMAL PROTECTION / MILD STRENGTHENING (6-12 Weeks Post-OP)

- A/AAROM No Limit FE (forward elevation in the scapular plane)
- A/AAROM No Limit ER (external rotation)
- 10-12 weeks, AIAA/PROM to improve ER with arm in **45 degree** abduction.
- AROM all directions below horizontal, light resisted motions in all planes.
- AROM activities to restore flexion, IR, horiz ADD as tolerated.
- Deltoid, Rotator Cuff isometrics progressing to isotonics.

- ❑ PRE's for scapular muscles, latissimus, biceps, triceps.
- ❑ PRE's work rotators in isolation (use modified neutral).
- ❑ Emphasize **posterior cuff, latissimus, and scapular muscle** strengthening, stressing eccentrics.
- ❑ Utilize **exercise arcs** that protect anterior and posterior capsule from stress during PRE's.
- ❑ Keep **all** strength exercises below the **horizontal plane** in this phase.

#### **Phase IV STRENGTHENING (12-16 Weeks Post-Op)**

- ❑ CRITERIA:
  - 1) Pain-free AROM
  - 2) Pain-free with manual muscle test
  - 3) Progress by response to treatment
- ❑ AROM activities to restore full ROM.
- ❑ Restore scapulohumeral rhythm.
- ❑ Joint mobilization.
- ❑ Aggressive scapular stabilization and eccentric strengthening program.
- ❑ Initiate isotonic shoulder strengthening exercises including: side lying ER, prone arm raises at 0, 90, 120 degrees, elevation in the plane of the scapula with IR and ER, lat pulldown close grip, and prone ER.
- ❑ Dynamic stabilization WB and NWB.
- ❑ PRE's for all upper quarter musculature (begin to integrate upper extremity patterns). Continue to emphasize eccentrics and glenohumeral stabilization.
- ❑ All PRE's are below the horizontal plane for non-throwers.
  - 1) Begin isokinetics.
  - 2) Begin muscle endurance activities (UBE).
    - High seat and low resistance
    - Must be able to do active shoulder flexion to 90 degrees without substitution
  - 3) Continue with agility exercises.
  - 4) Advanced functional exercises.
  - 5) Isokinetic test.
  - 6) Functional test assessment.
  - 7) Full return to sporting activities.