

Nikhil Verma, MD
Hailey Merk, MSM, MMS, PA-C
Lisa Raaf, MS, PA-C
1611 W. Harrison, Suite #300
Chicago, IL 60612
Vermapa@rushortho.com
Fax: 708-409-5179
www.sportssurgerychicago.com

MIDWEST
ORTHOPAEDICS
at RUSH



Post-Operative Rehabilitation Guidelines for Shoulder Arthroscopy Debridement

- 0-4 Weeks: Sling for Comfort (1-2 days) Then discontinue
Passive to Active shoulder ROM as tolerated
140° Forward Flexion
40° External Rotation with arm at side
Internal rotation behind back with gentle posterior capsule stretching
No rotation with arm in abduction until 4 wks
With distal clavicle excision, hold cross body adduction until 8wks.
Grip Strength, Elbow/Wrist/Hand ROM, Codmans
Avoid Abduction and 90/90 ER until 8wks
- 4-8 Weeks: Advance ROM as tolerated (Goals FF to 160°, ER to 60°)
Begin Isometric exercises
Progress deltoid isometrics
ER/IR (submaximal) at neutral
Advance to theraband as tolerated
- 8-12 Weeks: Advance to full, painless ROM
Continue strengthening as tolerated
Begin eccentrically resisted motion and closed chain activities
Only do strengthening 3times/wk to avoid rotator cuff tendonitis