

**Nikhil Verma, MD**  
**Lisa Raaf, MS, PA-C**  
**Lauren Rooney, MMS, PA-C**  
**Nathaniel Davidson, MS, PA-C**  
1611 W. Harrison, Suite #300  
Chicago, IL 60612  
Vermapa@rushortho.com  
Fax: 708-409-5179  
[www.sportssurgerychicago.com](http://www.sportssurgerychicago.com)



**MIDWEST  
ORTHOPAEDICS  
AT RUSH**



## **POSTOPERATIVE INSTRUCTIONS OPEN SHOULDER STABILIZATION (LATERJET)**

**\*\*Please note that the instructions provided below are general guidelines to be followed; however, any written or verbal instructions provided by Dr. Verma or either Physician Assistant supersede the instructions below and should be followed.**

### **DIET**

- Begin with clear liquids and light foods (jello, soups, etc.)
- Progress to your normal diet if you are not nauseated

### **WOUND CARE**

- Maintain your operative dressing, loosen bandage if swelling of the hand occurs
- It is normal for the shoulder to bleed and swell following surgery. If blood soaks through the bandage, do not become alarmed, reinforce with additional dressing
- Remove surgical dressing on the **7th post-operative day** – if minimal drainage is present, apply band-aids or a clean dressing over incisions and change daily.
- To avoid infection, keep surgical incisions clean and dry – you may shower by placing a plastic covering over the surgical site beginning the day after surgery. **NO** immersion of the arm.
- You can get your wound wet in the shower on the **7th post-operative day**. **NO** immersion in a bath until given approval by our office.

### **MEDICATIONS**

- Local anesthetics are injected into the wound and shoulder joint at the time of surgery. This will wear off within 8-12 hours and it is not uncommon for patients to encounter more pain on the first or second day after surgery when swelling peaks.
- Most patients will require some narcotic pain medication for a short period of time – this can be taken as per directions on the bottle.
- Common side effects of the pain medication are nausea, drowsiness, and constipation. To decrease the side effects take the medication with food. If constipation occurs, consider taking an over the counter laxative.
- If you are having problems with nausea and vomiting, contact the office to possibly have your medications changed.
- Do not drive a car or operate machinery while taking the narcotic medication or while in sling
- If you are having pain that is not being controlled by the pain medication prescribed, you may take an over the counter anti-inflammatory medication such as ibuprofen or naproxen in between doses of pain medication. This will help to decrease pain and decrease the amount of narcotic medication required. Please take as directed on the bottle.

- For 2 weeks following surgery take one 81mg aspirin twice daily to lower the risk of developing a blood clot after surgery. Please contact the office should severe distal arm pain occur or significant swelling of the distal arm and/or hand occur.
- 3-5 days of antibiotics post operatively will be prescribed to decrease chance of infection from the open incision.

### ACTIVITY

- You are to wear the sling placed at surgery for 4 weeks following surgery as described by Dr. Verma.
- When sleeping or resting, inclined positions (ie: reclining chair) and a pillow under the forearm for support may provide better comfort
- Do not engage in activities which increase pain/swelling. Unless otherwise instructed the arm should remain in the sling at all times.
- Avoid long periods of sitting or long distance traveling for 2 weeks.
- NO driving until instructed otherwise by physician, it is illegal to drive in a sling
- May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable

### IMMOBILIZER (if prescribed)

- Your sling is to be worn 4 weeks following surgery
- If there are 24 hours a day, the sling is to be worn 23.5 hours a day. Removal only for hygiene, dressing, and home exercises (as instructed by our office).

### ICE THERAPY

- Icing is very important in the initial post-operative period and should begin immediately after surgery.
- Use ice machine continuously for 45 minutes every 2 hours daily until your first post-operative visit. Care should be taken with icing to avoid frostbite to the skin.

### EXERCISE

- Begin exercises (active flexion/extension at elbow without resistance) 24 hours after surgery unless otherwise instructed.
- While maintaining your elbow by the side, begin elbow, hand, and wrist exercises immediately.
- Formal physical therapy (PT) typically begins after you are seen at your first post operative appointment 2 weeks after surgery. A prescription and protocol will be provided at your first post-op visit.

### EMERGENCIES\*\*

- Contact Dr. Verma's PA's at [Vermapa@rushortho.com](mailto:Vermapa@rushortho.com) if any of the following are present:
  - Painful swelling or numbness (note that some swelling and numbness is normal)
  - Unrelenting pain
  - Fever (over 101° - it is normal to have a low grade fever for the first day or two following surgery) or chills
  - Redness around incisions
  - Color change in distal arm and/or hand
  - Continuous drainage or bleeding from incision (a small amount of drainage is expected)
  - Difficulty breathing
  - Excessive nausea/vomiting
  - Calf pain
- If you have an emergency **after office hours** or on the weekend, contact the office at 312-432-2390 and you will be connected to our pager service. This will connect you with the Physician on call. You can also call Rush University Medical Center at **312-942-5000** and ask for the operator to page the orthopedic resident on call.
- If you have an emergency that requires immediate attention proceed to the nearest emergency room.

### FOLLOW-UP CARE/QUESTIONS

- If you do not already have a post-operative appointment scheduled, please contact 708-236-2701.
- Typically the first post-operative appointment following surgery is 10-14 days following surgery
- The first post operative appointment will be with one of the Physician Assistants. They will assess the wound, go over post operative protocol, and answer any questions you may have regarding the procedure
- If you have any further questions please contact the PAs directly at [Vermapa@rushortho.com](mailto:Vermapa@rushortho.com)